



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2018

---

## **The use of levosimendan in cardiac surgery: an update after the LEVO-CTS, CHEETAH and LICORN trials in the light of clinical practice**

Guarracino, Fabio ; Heringlake, Matthias ; Cholley, Bernard ; Bettex, Dominique ; Bouchez, Stefaan ; Lomivorotov, Vladimir V ; Rajek, Angela ; Kivikko, Matti ; Pollesello, Piero

DOI: <https://doi.org/10.1097/FJC.0000000000000551>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-142247>

Journal Article

Accepted Version

Originally published at:

Guarracino, Fabio; Heringlake, Matthias; Cholley, Bernard; Bettex, Dominique; Bouchez, Stefaan; Lomivorotov, Vladimir V; Rajek, Angela; Kivikko, Matti; Pollesello, Piero (2018). The use of levosimendan in cardiac surgery: an update after the LEVO-CTS, CHEETAH and LICORN trials in the light of clinical practice. *Journal of Cardiovascular Pharmacology*, 71(1):1-9.

DOI: <https://doi.org/10.1097/FJC.0000000000000551>

**The use of levosimendan in cardiac surgery: an update after the LEVO-CTS,  
CHEETAH and LICORN trials in the light of clinical practice**

*Fabio Guarracino, MD, PhD,\* Matthias Heringlake, MD, PhD,†  
Bernard Cholley, MD, PhD, ‡§ Dominique Bettex, MD, PhD,¶ Stefaan Bouchez, MD, PhD, ||  
Vladimir V.Lomivorotov, MD, PhD,\*\* Angela Rajek, MD,†† Matti Kivikko, MD, PhD, ‡‡§§  
and Piero Pollesello, PhD‡‡*

\*Dipartimento di Anesthesia e Terapie Intensive, Azienda Ospedaliero-Universitaria Pisana, Pisa, Italy; †Klinik für Anästhesiologie und Intensivmedizin, Universitätsklinikum Schleswig-Holstein, Lübeck, Germany; ‡Department of Anesthesiology and Critical Care Medicine, Hôpital Européen Georges Pompidou, Paris, France; §Université Paris Descartes, Sorbonne Paris Cité; Paris, France; ¶Cardiac Anaesthesia, University Hospital, Zurich, Switzerland; ||Department of Anesthesiology, University Hospital, Ghent, Belgium; \*\*Department of Anesthesiology, E.Meshalkin National Medical Research Center, Novosibirsk, Russia; ††Klinische Abteilung für Herz-Thorax-Gefäßchirurgische Anästhesie und Intensivmedizin, Medical University of Vienna, Vienna, Austria; ‡‡Critical Care Proprietary Products, Orion Pharma, Espoo, Finland; §§Department of Cardiology S7, Jorvi Hospital, Espoo, Finland.

**Reprints:** Piero Pollesello, PhD, Adj.Prof., FESC, Critical Care Proprietary Products, Orion Pharma, P.O. Box 65, 02101 Espoo, Finland. Tel: +358509664191; E-mail: [piero.pollesello@orionpharma.com](mailto:piero.pollesello@orionpharma.com)

This is an open access article distributed under the terms of the Creative Commons Attribution License 4.0 (CC BY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### **Abstract**

Levosimendan is a calcium sensitizer and ATP-dependent potassium channel opener which exerts sustained hemodynamic, symptomatic and organ-protective effects. It is registered for the treatment of acute heart failure, and when inotropic support is considered appropriate. In the past fifteen years, levosimendan has been widely used in clinical practice, and has also been tested in clinical trials to stabilize at-risk patients undergoing cardiac surgery. Recently, three randomized, placebo-controlled, multicenter studies (LICORN, CHEETAH and LEVO-CTS) have been published reporting on the peri-operative use of levosimendan in patients with compromised cardiac ventricular function. Taken together, many smaller trials conducted in the past suggested beneficial outcomes with levosimendan in peri-operative settings. In contrast, the latest three studies were neutral or inconclusive. In order to understand the reasons for such dissimilarity, a group of experts from Austria, Belgium, Finland, France, Germany, Italy, Switzerland, and Russia, including investigators from the three most recent studies, met to discuss the study results in the light of both the previous literature and current clinical practice. Despite the fact that the null hypothesis could not be

ruled out in the recent multicenter trials, we conclude that levosimendan can still be viewed as a safe and effective inodilator in cardiac surgery.

**Short title:** latest trials on levosimendan in cardiac surgery

**Key words:** cardiac surgery; clinical trials; levosimendan; systematic review; opinion paper.

## 1. Introduction

Peri-operative mortality is reported to be as low as 1–4% in the general elective surgery population.<sup>1</sup> However, in patients with postoperative low cardiac output syndrome (LCOS), mortality is considerably higher.<sup>2</sup> In addition to higher mortality, LCOS predisposes patients to postoperative myocardial injury, renal failure, and prolonged intensive care unit and hospital stay.<sup>3</sup> Several baseline factors, such as preoperatively reduced left (and/or right) ventricular function or recent myocardial infarction predispose patients to LCOS.<sup>4</sup> The type of surgery also affects the postoperative risk profile; coronary artery bypass grafting (CABG) alone has a more benign outcome than, for example, the combination of CABG and a valve replacement.<sup>5</sup>

LCOS is managed with inotropic agents and/or mechanical cardiac assist devices such as an intra-aortic balloon pump (IABP). Even so, short-term mortality is greatly elevated versus non-LCOS comparators.<sup>6</sup> Moreover, the inotropic agents traditionally used in this setting have conspicuous adverse effects or incompletely defined safety profiles.<sup>7</sup>

Levosimendan is a calcium sensitizer and ATP-dependent potassium channel opener with positive inotropic, vasodilatory and cardioprotective properties.<sup>8</sup> The drug binds to cardiac troponin C in a calcium-dependent manner<sup>9,10</sup> which mediates the positive inotropic effect by increasing the calcium sensitivity of myocytes. The vasodilatory effect is due to the opening of ATP-sensitive potassium channels in vascular smooth muscle resulting in its relaxation. By opening mitochondrial ATP-sensitive potassium channels in cardiomyocytes the drug also exerts a cardioprotective effect. In addition, inhibition of phosphodiesterase III by levosimendan has been also proposed to have a role in its pharmacodynamics effects.<sup>11</sup>

Levosimendan has been in clinical use for 15 years. In addition to its original indication for acutely decompensated heart failure, it has also been used to stabilize patients undergoing cardiac surgery. Abundant literature from exploratory studies supports the rationale for its use in this indication,<sup>12</sup> and this is also supported by its benign effect on kidney function.<sup>13</sup>

Recently, three randomized, placebo-controlled, multicenter studies were published on the peri-operative use of levosimendan: LICORN,<sup>14</sup> CHEETAH,<sup>15</sup> and LEVO-CTS.<sup>16</sup> In contrast to the many preceding smaller trials which, either individually or as a whole, produced a promising image of levosimendan in peri-operative settings, these latest three studies were either neutral or inconclusive.

A group of experts from eight European countries (Austria, Belgium, Finland, France, Germany, Italy, Switzerland, and Russia), including investigators from the three most recent studies on the pre-, peri-, and postoperative use of levosimendan, met on 20 April 2017 in occasion of the EACTA annual congress in Berlin, Germany, to discuss the recent study results in the light of both the previous literature and current clinical practice. The present paper was created from the proceedings of that discussion.

## 2. Previous relevant studies on the use of levosimendan in cardiac surgery

Levosimendan has been studied in >40 clinical trials in cardiac surgery. Earlier studies suggested that it could prevent the development of LCOS and be effective in treating postoperative LCOS (see **BOX 1**). The level of proof, however, remained low despite a meta-analysis that suggested a survival benefit in patients with low pre-operative ejection fraction.<sup>17</sup> Indications of renal-protective effects in this setting have also been reported in retrospective analyses.<sup>18,19</sup>

The individual and aggregate findings of the 14 studies in cardiac surgery patients with low left ventricular ejection fraction (LVEF) examined by Harrison et al.<sup>17</sup> in their meta-analysis are reported in **Figure 1**, and the results of the contributing studies are summarized briefly in **Table 1**.<sup>20-33</sup>

In total, Harrison and coworkers took into consideration data from 1155 patients (84 deaths) and the overall effect of levosimendan vs comparator was significant ( $p=0.008$ ). Visual inspection of the funnel plot for the primary outcome of mortality was not suggestive of significant publication bias. However, removal of either of the two studies by Levin and coworkers<sup>25,26</sup> made the overall estimated effect of levosimendan on mortality insignificant.

For the sake of completeness, we report that, in addition to the 14 papers considered by Harrison et al., other relevant studies conducted on levosimendan in surgical patients are those by Severi et al.<sup>34</sup> and Lomivorotov et al.<sup>35</sup> on levosimendan versus IABP; the trials by Baysal et al.<sup>36</sup> and Erb et al.<sup>37</sup> on renal outcome and organ dysfunction, respectively; and the randomized pilot study on prophylactic use of levosimendan by Anastasiadis et al.<sup>38</sup>

### 3. The three most recent studies

Three larger clinical studies have recently been conducted with levosimendan in patients undergoing cardiac surgery. All three were randomized, placebo-controlled, multicenter studies. The LICORN and CHEETAH studies were investigator-initiated studies, whereas LEVO-CTS was a phase 3 regulatory study (see **BOX 2**). Broad outlines of the study designs and their primary findings are given below.

#### 3.1. LICORN

The LICORN trial (Levosimendan on Low Cardiac Output Syndrome in Patients With Low Ejection Fraction Undergoing Coronary Artery Bypass Grafting With Cardiopulmonary Bypass; NCT02184819) assessed the efficacy of a pre-operative infusion of levosimendan in reducing postoperative LCOS in patients with poor LVEF undergoing CABG.<sup>14</sup> A cohort of 336 patients with LVEF  $\leq 40\%$  undergoing CABG was recruited from 13 French hospitals. The study drug was started after induction of anesthesia and infused over a period of 24 h at a rate of 0.1  $\mu\text{g/kg/min}$ . Postoperative LCOS was evaluated using a composite criterion comprising (1) need for inotropic agents beyond 48 h following discontinuation of the study drug; (2) need for postoperative mechanical assist devices or failure to wean from these devices when inserted pre-operatively; and (3) need for renal replacement therapy.

The primary endpoint was observed in 87/167 patients (52%) in the levosimendan group, compared with 101/168 (61%) in the placebo group (absolute risk reduction -7%, 95% confidence interval [CI]: -17% to +3%,  $p=0.15$ ). Of the secondary endpoints, the duration of catecholamine treatment was shorter in the levosimendan group:  $3.2 \pm 3.6$  versus  $4.1 \pm 4.3$  days ( $p=0.021$ ). However, no adjustment was made for multiple comparisons, and this result

should be considered exploratory. There were no statistically significant inter-group differences in mortality or length of ICU stay.

The lack of statistical significance in the composite primary endpoint was likely contributed to by the fact that the observed event rate in the placebo arm was lower than that anticipated (61% vs 65%); it was anticipated also that the prevalence rate would be reduced to 50% in the levosimendan group, but the prevalence actually observed was 52% in the intention-to-treat population (vs. 51% in the per-protocol population). LICORN was powered according to an expectation of an absolute risk reduction of 15%. The point estimate actually recorded was 7% and favored levosimendan but the 95% CI included a reduction of 17% (range -17% to +3%). The observed effect was less than that anticipated by the study hypothesis; however, the study was underpowered to definitely exclude a meaningful beneficial effect of levosimendan on the primary composite outcome.

### **3.2. CHEETAH**

In the CHEETAH (Levosimendan to Reduce Mortality in High Risk Cardiac Surgery Patients; NCT00994825) trial, levosimendan or placebo was administered to cardiac surgery patients, who, according to predefined criteria, developed postoperative LCOS.<sup>15</sup> In total, 1000 patients were scheduled to be included and the primary endpoint was 30-day mortality. The study was performed in 14 centers in Italy, Russia and Brazil but was stopped for futility after 506 patients had been enrolled. A total of 248 patients received levosimendan and 258 received placebo. The mean infusion rate and duration of levosimendan were 0.07 µg/kg/min for 33 h, and the median EF was 50% in both groups, with 11% of patients having an EF of < 25%. There was no significant difference in 30-day mortality between the levosimendan and placebo groups: 32 patients (12.9%) versus 33 (12.8%); absolute risk difference 0.1



percentage points; 95% CI –5.7 to +5.9 percentage points; P=0.97). There were no significant between-group differences in other endpoints and no difference in the rates of adverse events (hypotension or arrhythmias).

It should be noted that, in the report of the CHEETAH study, preparation of the study drug was described as follows:<sup>15</sup> “levosimendan was diluted as 12.5 mg in 100 ml of 5% glucose”. This is at variance with the summary of product characteristics guidance, according to which one vial of Simdax (12.5 mg levosimendan concentrate for i.v.) should be diluted in at least 250 mL of 5% glucose solution (1:50). There is a risk of precipitation if smaller diluent volumes are used and this exposes the patient to unpredictable dosing (i.e. receipt of less than the intended dose).

### **3.3 LEVO-CTS**

This study (Levosimendan in Patients with Left Ventricular Systolic Dysfunction Undergoing Cardiac Surgery Requiring Cardiopulmonary Bypass; NCT02025621) was a Phase III clinical trial sponsored by TENAX Therapeutics Inc., run by Duke University (Durham, NC, USA) and configured to support a marketing authorization in the U.S. and Canada for levosimendan.<sup>16</sup> The study design and objectives were discussed beforehand and agreed with the U.S. Food and Drug Administration (FDA).

The study population consisted of 882 patients with low pre-operative LVEF (EF<35%) undergoing scheduled or urgent cardiac surgery (CABG and/or mitral valve surgery with or without involvement of other valves). All patients were considered at risk of developing postoperative LCOS. Levosimendan (0.2 µg/kg/min for 60 min, followed by 0.1

µg/kg/min for 23 h) or placebo was started at the induction of anesthesia to assess whether the drug would decrease the development of LCOS and its detrimental consequences.

The study, conducted at 70 sites in Canada and the U.S., demonstrated no statistically robust treatment effect on the composite primary endpoint of death, peri-operative myocardial infarction and need for renal replacement therapy or a mechanical ventricular assist device. However, there were fewer deaths in the levosimendan group: 20/428 (4.7%) versus 30/421 (7.1%), odds ratio 0.64, 95% CI 0.37–1.13 ( $p=0.12$ ). In addition, the levosimendan-treated patients experienced statistically significantly fewer LCOS events (78 vs 108;  $p=0.007$ ) and needed less inotropic support at or beyond 24 h after initiation of infusion (235 vs. 264;  $p=0.02$ ). Cardiac index also improved more in levosimendan-treated patients ( $2.9\pm0.6$  vs  $2.7\pm0.7$  L/min/m<sup>2</sup>;  $p<0.001$ ). Hypotension and atrial arrhythmias were recorded as adverse events with similar frequency in both study groups.

The LEVO-CTS investigators are conducting post-hoc analyses in relevant predetermined sub-settings (e.g. CABG with or without accompanying valve surgery). So far, they have shown data suggesting that in those patients in whom only CABG was performed, mortality was significantly lower in the levosimendan group: 6/284 (2.1%) versus 22/279 (7.9%), hazard ratio 0.259, 95% CI 0.105–0.640 ( $p=0.0016$ ) (**Figure 2**).

## 4. Interpretation of the data

### 4.1. Efficacy

The hypotheses tested in all three studies were not affirmed, and the primary endpoints did not differ significantly between the levosimendan and control arms.

It must be noted that the endpoints used in LEVO-CTS and LICORN were experimental ones. Similar endpoints have not been used in earlier studies. The LEVO-CTS primary endpoints were agreed with the FDA, which required the inclusion of clinical events. This notwithstanding, encouraging evidence of efficacy emerged from LEVO-CTS: the lower incidence of LCOS, lesser need for inotropic support by catecholamines, and improvement in CI indicate that levosimendan exhibited efficacy. In CHEETAH, only suggestive signs of improved hemodynamics were noted but the dose of levosimendan in that study was smaller than those in LEVO-CTS or LICORN or in previous studies in cardiac surgery. The subgroup data from LEVO-CTS (the supplementary material of the publication is available via <http://www.nejm.org/doi/full/10.1056/NEJMoa1616218>) suggest that levosimendan may be more effective in patients in whom only the CABG procedure is performed (and ineffective in valve replacement patients). Also, in line with earlier data, patients with the lowest EF before surgery may benefit most from the treatment. However, this was not observed in the LICORN study, in which no statistically significant difference was found with respect to the primary endpoint between patients with LVEF<30% or between 30% and 40%.

Although the duration of treatment with inotropic agents was an outcome variable, the LICORN and CHEETAH protocols do not give details of hemodynamic monitoring and specifically when inotrope treatment should be stopped. With regard to the limited efficacy of levosimendan, the regimen used in the trials can be questioned. Both LEVO-CTS and LICORN used a 0.1 µg/kg/min infusion for 24 h without bolus, in conformity with recent recommendations by experts in cardiovascular anesthesia.<sup>12</sup> Several previous studies have used infusion rates of 0.2 µg/kg/min with variable bolus doses. Higher doses may have produced greater hemodynamic effects, but at the expense of a more potent vasodilatation and consequent hypotension

In the LICORN trial, inodilators and inopressors were analyzed together as ‘catecholamines’ irrespective of their predominant hemodynamic effect and no doses for the respective drugs were reported. Consequently it cannot be ruled out that patients with a severe postoperative LCOS receiving high doses of epinephrine and milrinone, and those needing only small doses of norepinephrine on the second day after surgery were both classified as ‘catecholamine-dependent’.

One additional observation regarding LEVO-CTS and LICORN is that levosimendan therapy was started very shortly before surgery; accordingly, there was only a short time during which levosimendan could exert any preconditioning effect. In some previous studies, and in clinical practice, levosimendan has been administered for up to 24 h before the start of surgery.<sup>26,30</sup>

Finally, in the CHEETAH trial the majority of patients received a relatively low dose of levosimendan whilst already being treated with high doses of epinephrine and dobutamine. Pre-treatment with beta-mimetic drugs has been shown to reduce the inotropic effect of levosimendan *in vitro*<sup>39</sup> and thereby may also reduce its benefits *in vivo*, as shown by Bonios et al.<sup>40</sup> in a trial comparing the event-free survival of patients treated with levosimendan, dobutamine, or their combination.

#### **4.2. Safety**

Safety was not identified as a concern in any of these three studies: there was no significant excess of arrhythmias or hypotension and no increase in mortality in levosimendan-treated patients. In fact, mortality was numerically lower in levosimendan-treated patients in LEVO-CTS.<sup>16</sup>

These findings are fully consistent with the data on safety and adverse events reported in most of the previous studies (as collected in a systematic review and meta-analysis by Landoni et al.<sup>41</sup>) and confirm levosimendan as the safest agent among the family of inotropes and inodilators which include, among others, dobutamine and milrinone.<sup>42</sup> When an inotrope is needed, the safety profile of the chosen agent should be an important selection criterion.

In addition, levosimendan has a unique mechanism of action and pharmacokinetics. The sensitization of myofilaments to calcium supposes that the inotropic effects occurs without (or with minimal) increase in myocardial oxygen consumption. The prolonged effect which lasts for several days,<sup>43,44</sup> contrasts with the on-off action of dobutamine. These specificities may prove useful in selected patients and are a major reason to keep levosimendan in the armamentarium of physicians in charge of patients with cardiac dysfunction.

## 5. Discussion

While the many smaller trials conducted in the past produced as a whole a promising image of levosimendan in peri-operative settings, the data from the latest (and larger) three studies did not support the hypotheses tested.

Instead of advocating the ‘small-study effect’, i.e. the trend for smaller studies to show larger treatment effects,<sup>45</sup> every study should be evaluated fairly, as large studies can be imprecise, just as small ones can be precise.<sup>46</sup>

Indeed, in small monocentric trials focused on a few endpoints in very specific clinical settings, the researchers are in a more controlled situation than in a multicenter trial in the field. In addition, in clinical settings where there are local variations in therapeutic approaches

and tailored strategies (such as, for example, in mitral valve surgery<sup>47</sup>), multicenter trials add, by definition, statistical noise to many endpoints: variations in pharmacologic and non-pharmacologic parameters have the potential to impair the statistical power and obscure meaningful effects. Multicenter studies in fields such as perioperative LCOS, where multiple sources of heterogeneity exist (e.g. symptoms, etiologies, comorbidities, co-medications and center-specific treatment practices), encounter such problems.

All the above notwithstanding, we have noticed that studies of all sizes on levosimendan have produced some common and consistent findings in terms of safety. Levosimendan is safe and well tolerated in patients undergoing cardiac surgery with cardiopulmonary bypass who have low LVEF and are at risk of the development of postoperative LCOS. This safety finding, and especially the lack of any deterioration in survival, is particularly noteworthy given that many of the patients in all three trials had already been treated with a range of other pressor and/or inotropic drugs. The non-attainment of the study hypotheses in these three recent trials does not rule out the fact that levosimendan might be effective in selected patients undergoing cardiac surgery. The LEVO-CTS study, as the largest of these trials, confirms that a prophylactic infusion of levosimendan started immediately before surgery reduces LCOS in a heterogeneous population of cardiac surgery patients with reduced LVEF. The post-hoc analyses further suggest that this drug may be especially useful in patients undergoing CABG with reduced LVEF, but not in those undergoing a valve surgery. A recent meta-analysis, including also the latest trials, confirms that levosimendan had a significant effect on mortality only when used in case of severe perioperative cardiovascular dysfunction ( $\text{LVEF} \leq 30\%$ ) in patients receiving cardiac surgery.<sup>48</sup> No comparable data are available for any other drugs with inotropic properties; on the contrary, traditional inotropes are considered to have detrimental effects on outcome.<sup>42</sup>

In addition, levosimendan has been shown to reduce elevated right-sided pressures in various clinical situations.<sup>49-51</sup> Preoperative administration of levosimendan decreased pulmonary artery pressure significantly in patients with right ventricular dysfunction and pulmonary hypertension.<sup>52</sup> As pulmonary hypertension is an important prognostic factor in cardiac surgery associated with increased morbidity and mortality,<sup>53</sup> levosimendan's efficacy could be pronounced in this subgroup and we warmly suggest to explore this setting.

Finally, it must be registered that in many studies, including relatively large regulatory clinical trials, levosimendan was administered in addition to standard of care (i.e. other vasoactive drugs) according to prevailing practice at individual study centers. It would be instructive to perform a post-hoc analysis of those data to explore whether combinations of levosimendan with dobutamine or norepinephrine are beneficial.

## 6. Conclusions

Taking all the available data into consideration, including the experience of the three most recent studies, our conclusion is that levosimendan is a safe and effective agent for the treatment of patients undergoing cardiac surgery and in need of inotropic support (see **BOX 3**). However, the magnitude of effect of this agent is not as large as previously thought and three large multi-center trials could not rule out their null hypothesis. For this reason levosimendan cannot be at the moment recommended for routine use in all cardiac surgery settings. Further in-depth assessment of the utility of levosimendan will require additional trials in closely defined patient populations with study designs that mitigate, to the fullest extent possible, any influence of methodological variations in patient management.

## **Acknowledgments**

Hughes associates (Oxford, UK) provided editorial assistance in the preparation of this article.

## **Declarations**

Levosimendan is approved in 61 world countries for treatment of acute decompensated heart failure, and when inotropic support is considered appropriate. In the US the drug does not currently have a market authorization by FDA, but is under development for prevention of Low cardiac Output Syndrome in patients at risk who undergo a cardiac operation.

This project did not receive any financial support, apart from logistical expenses relating to the organization of the consensus meeting at the EACTA annual congress in Berlin, Germany, on April 20 2017, which were covered by Orion Pharma. The attendees, who were identified on the basis of their experience with peri-operative use of levosimendan, as documented in the peer-reviewed literature, did not receive honoraria for this specific activity. MH reports honoraria for scientific advice and lectures from Orion Pharma and Tenax Therapeutics. MK and PP are full-time employee of Orion Pharma, where levosimendan was discovered and developed.



## References

1. Pearse RM, Moreno RP, Bauer P, et al.; European Surgical Outcomes Study (EuSOS) group for the Trials groups of the European Society of Intensive Care Medicine and the European Society of Anaesthesiology. Mortality after surgery in Europe: a 7 day cohort study. *Lancet*. 2012;380:1059-1065.
2. Coulson TG, Bailey M, Reid CM, et al. The association between peri-operative acute risk change (ARC) and long-term survival after cardiac surgery. *Anaesthesia*. 2017, Jul 13. [epub ahead of print, doi: 10.1111/anae.13967]
3. Pérez Vela JL, Jiménez Rivera JJ, Alcalá Llorente MÁ, et al.; en representación del Grupo ESBAGA. Low cardiac output syndrome in the postoperative period of cardiac surgery. Profile, differences in clinical course and prognosis. The ESBAGA study. *Med Intensiva*. 2017, Jul 20. [Epub ahead of print, doi: 10.1016/j.medin.2017.05.009]
4. Algarni KD, Maganti M, Yau TM. Predictors of low cardiac output syndrome after isolated coronary artery bypass surgery: trends over 20 years. *Ann Thorac Surg*. 2011;92(5):1678-1684.
5. Hannan EL, Wu C, Bennett EV, Carlson RE, et al. Risk index for predicting in-hospital mortality for cardiac valve surgery. *Ann Thorac Surg*. 2007;83(3):921–929.
6. Hillis LD, Smith PK, Anderson JL, et al.. 2011 ACCF/AHA Guideline for Coronary Artery Bypass Graft Surgery: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *Circulation*. 2011;124(23):e652-735.
7. Nielsen DV, Hansen MK, Johnsen SP, et al. Health outcomes with and without use of inotropic therapy in cardiac surgery: results of a propensity score-matched analysis. *Anesthesiology*. 2014;120(5):1098-1108.

8. Papp Z, Édes I, Fruhwald S, et al. Levosimendan: molecular mechanisms and clinical implications: consensus of experts on the mechanisms of action of levosimendan. *Int J Cardiol.* 2012;159(2):82-87
9. Pääkkönen K, Annala A, Sorsa T, et al. Solution structure and main chain dynamics of the regulatory domain (residues 1–91) of human cardiac troponin C. *J.Biol. Chem.* 1998;273(25):15633-15638
10. Sorsa T, Pollesello P, P Permi P, et al. Interaction of levosimendan with cardiac troponin C in the presence of cardiac troponin I peptides. *J. Mol. Cell. Cardiol.* 2003;35(9):1055-1061
11. Endoh M. *J Cardiovasc Pharmacol.* 2002;40:323-338.
12. Toller W, Heringlake M, Guarracino F, et al. Preoperative and perioperative use of levosimendan in cardiac surgery: European expert opinion. *Int J Cardiol.* 2015;184:323-336.
13. Yilmaz MB, Grossini E, Silva Cardoso JC, et al. Renal effects of levosimendan: a consensus report. *Cardiovasc Drugs Ther.* 2013;27(6):581-590
14. Cholley B, Caruba T, Grosjean S, et al. Effect of Levosimendan on Low Cardiac Output Syndrome in Patients With Low Ejection Fraction Undergoing Coronary Artery Bypass Grafting With Cardiopulmonary Bypass: The LICORN Randomized Clinical Trial. *JAMA.* 2017;318:548-556
15. Landoni G, Lomivorotov VV, Alvaro G, et al.; CHEETAH Study Group. Levosimendan for Hemodynamic Support after Cardiac Surgery. *N Engl J Med.* 2017;376(21):2021-2031.

16. Mehta RH, Leimberger JD, van Diepen S, et al.; LEVO-CTS Investigators.  
Levosimendan in Patients with Left Ventricular Dysfunction Undergoing Cardiac Surgery. *N Engl J Med*. 2017;376(21):2032-2042.
17. Harrison RH, Hasselblad V, Mehta RH, et al. Effect of levosimendan on survival and adverse events after cardiac surgery: a meta-analysis. *J Cardiothorac Vasc Anesth*. 2013;27:1224-1232.
18. Grieshaber P, Lipp S, Arnold A, et al. Impact of prophylactic administration of Levosimendan on short-term and long-term outcome in high-risk patients with severely reduced left-ventricular ejection fraction undergoing cardiac surgery - a retrospective analysis. *J Cardiothorac Surg*. 2016;11(1):162.
19. Balzer F, Treskatsch S, Spies C, et al. Early administration of levosimendan is associated with improved kidney function after cardiac surgery - a retrospective analysis. *J Cardiothorac Surg*. 2014;9:167.
20. Al-Shawaf E, Ayed A, Vislocky I, et al. Levosimendan or milrinone in the type 2 diabetic patient with low ejection fraction undergoing elective coronary artery surgery. *J Cardiothorac Vasc Anesth*. 2006;20(3):353-357.
21. Alvarez J, Taboada M, Rodríguez J, et al. Hemodynamic effects of levosimendan following cardiac surgery. *Rev Esp Anesthesiol Reanim*. 2005;52(7):389-394.
22. Alvarez J, Bouzada M, Fernández AL, et al. Hemodynamic effects of levosimendan compared with dobutamine in patients with low cardiac output after cardiac surgery. *Rev Esp Cardiol*. 2006;59(4):338-345.
23. De Hert SG, Lorsomradee S, Cromheecke S, et al. The effects of levosimendan in cardiac surgery patients with poor left ventricular function. *Anesth Analg*. 2007;104(4):766-773.

24. Eriksson HI, Jalonon JR, Heikkinen LO, et al. Levosimendan facilitates weaning from cardiopulmonary bypass in patients undergoing coronary artery bypass grafting with impaired left ventricular function. *Ann Thorac Surg.* 2009;87(2):448-454.
25. Levin R, Porcile R, Salvagio F, et al. Levosimendan reduces mortality in post-operative low cardiac output syndrome after coronary surgery. *Circulation.* 2009;120(suppl. 18):S987-S988.
26. Levin R, Degrange M, Del Mazo C, et al. Preoperative levosimendan decreases mortality and the development of low cardiac output in high-risk patients with severe left ventricular dysfunction undergoing coronary artery bypass grafting with cardiopulmonary bypass. *Exp Clin Cardiol.* 2012;17(3):125-130.
27. Lomivorotov VV, Cherniavskiy AM, Boboshko VA, et al. Levosimendan vs. intra-aortic balloon pump in high-risk cardiac surgery. *Asian Cardiovasc Thorac Ann.* 2011;19(2):154-159.
28. Järvelä K, Maaranen P, Sisto T, et al. Levosimendan in aortic valve surgery: cardiac performance and recovery. *J Cardiothorac Vasc Anesth.* 2008;22(5):693-698
29. Lahtinen P, Pitkänen O, Pölönen P, et al. Levosimendan reduces heart failure after cardiac surgery: a prospective, randomized, placebo-controlled trial. *Crit Care Med.* 2011;39(10):2263-2270
30. Leppikangas H, Järvelä K, Sisto T, et al. Preoperative levosimendan infusion in combined aortic valve and coronary bypass surgery. *Br J Anaesth.* 2011;106(3):298-304
31. Momeni M, Rubay J, Matta A, et al. Levosimendan in congenital cardiac surgery: a randomized, double-blind clinical trial. *J Cardiothorac Vasc Anesth.* 2011;25(3):419-424
32. Tritapepe L, De Santis V, Vitale D, et al. Preconditioning effects of levosimendan in coronary artery bypass grafting--a pilot study. *Br J Anaesth.* 2006;96(6):694-700

33. Tritapepe L, De Santis V, Vitale D, et al. Levosimendan pre-treatment improves outcomes in patients undergoing coronary artery bypass graft surgery. *Br J Anaesth.* 2009;102(2):198-204
34. Severi L, Lappa A, Landoni G, et al. Levosimendan versus intra-aortic balloon pump in high-risk cardiac surgery patients. *J Cardiothorac Vasc Anesth.* 2011;25(4):632-636.
35. Lomivorotov VV, Boboshko VA, Efremov SM, et al. Levosimendan versus an intra-aortic balloon pump in high-risk cardiac patients. *J Cardiothorac Vasc Anesth.* 2012;26(4):596-603.
36. Baysal A, Yanartas M, Dogukan M, et al. Levosimendan improves renal outcome in cardiac surgery: a randomized trial. *J Cardiothorac Vasc Anesth.* 2014;28(3):586-594.
37. Erb J, Beutlhauser T, Feldheiser A, et al. Influence of levosimendan on organ dysfunction in patients with severely reduced left ventricular function undergoing cardiac surgery. *J Int Med Res.* 2014;42(3):750-764.
38. Anastasiadis K, Antonitsis P, Vranis K, et al. Effectiveness of prophylactic levosimendan in patients with impaired left ventricular function undergoing coronary artery bypass grafting: a randomized pilot study. *Interact Cardiovasc Thorac Surg.* 2016;23(5):740-747.
39. Haikala H, Kaheinen P, Levijoki J, et al. The role of cAMP- and cGMP-dependent protein kinases in the cardiac actions of the new calcium sensitizer, levosimendan. *Cardiovasc Res.* 1997;34(3):536-546.
40. Bonios MJ, Terrovitis JV, Drakos SG, et al. Comparison of three different regimens of intermittent inotrope infusions for end stage heart failure. *Int J Cardiol.* 2012;159(3):225-229.

41. Landoni G, Biondi-Zoccai G, Greco M, et al. Effects of levosimendan on mortality and hospitalization. A meta-analysis of randomized controlled studies. *Crit Care Med*. 2012;40(2):634-646
42. Greco T, Calabrò MG, Covello RD, et al. A Bayesian network meta-analysis on the effect of inodilatory agents on mortality. *Br J Anaesth*. 2015;114(5):746-756.
43. De Hert SG, Lorsomradee S, Cromheecke S, et al. The effects of levosimendan in cardiac surgery patients with poor left ventricular function. *Anesth Analg*. 2007;104(4):766-773.
44. Kivikko M, Lehtonen L, Colucci WS. Sustained hemodynamic effects of intravenous levosimendan. *Circulation*. 2002;107:81-86.
45. Sterne JA, Gavaghan D, Egger M. Publication and related bias in meta-analysis: power of statistical tests and prevalence in the literature. *J.Clin.Epidemiol*. 2000; 53:1119–1129.
46. Choi SW and Lam DM. Funnels for publication bias - have we lost the plot? *Anaesthesia*. 2016;71(3):338-341.
47. Andalib A, Chetrit M, Eberg M, et al. A Systematic Review and Meta-Analysis of Outcomes Following Mitral Valve Surgery in Patients with Significant Functional Mitral Regurgitation and Left Ventricular Dysfunction. *J Heart Valve Dis*. 2016;25(6):696-707.
48. Lee C-T, Lin Y-C, Yeh Y-C, et al. Effects of levosimendan for perioperative cardiovascular dysfunction in patients receiving cardiac surgery: a meta-analysis with trial sequential analysis. *Intensive Care Med* 2017 [published online Sept 7, 2017] DOI: 10.1007/s00134-017-4927-5.
49. Kleber FX, Bollmann T, Borst MM, et al. Repetitive dosing of intravenous levosimendan improves pulmonary hemodynamics in patients with pulmonary hypertension: results of a pilot study. *J Clin Pharmacol*. 2009;49(1):109-115.

50. Ebade AA, Khalil MA, Mohamed AK. Levosimendan is superior to dobutamine as an inodilator in the treatment of pulmonary hypertension for children undergoing cardiac surgery. *J Anesth*. 2013;27(3):334-339.
51. Morelli A, Teboul JL, Maggiore SM, et al. Effects of levosimendan on right ventricular afterload in patients with acute respiratory distress syndrome: a pilot study. *Crit Care Med*. 2006;34(9):2287-2293.
52. Guerrero-Orriach JL, Ariza-Villanueva D, Florez-Vela A, et al. Cardiac, renal, and neurological benefits of preoperative levosimendan administration in patients with right ventricular dysfunction and pulmonary hypertension undergoing cardiac surgery: evaluation with two biomarkers neutrophil gelatinase-associated lipocalin and neuronal enolase. *Ther Clin Risk Manag*. 2016;12:623-630.
53. Denault A, Deschamps A, Tardif JC, et al. Pulmonary hypertension in cardiac surgery. *Curr Cardiol Rev*. 2010;6(1):1-14.

## Legends to the figures

**Figure 1.** Meta-analysis of data from 14 randomized controlled trials of peri-operative levosimendan in cardiac surgery patients (n=1155) indicates that levosimendan therapy is associated with reduced mortality, with the greatest benefit observed in patients with reduced LVEF. From Harrison and coworkers.<sup>17</sup>

**Figure 2.** Ninety-day mortality among patients in the LEVO-CTS trial in (A) the whole study (n=849) and (B) the subgroup of isolated CABG patients (n=563). In the latter, mortality was significantly lower in the levosimendan arm than in the placebo arm. From supplemental materials in Mehta and coworkers.<sup>16</sup>

**Figure 3.** Meta-analysis of clinical trials on levosimendan versus control in cardiac surgery patients with long term and thirty-day mortality as primary outcome: effects of levosimendan when used in case of severe perioperative cardiovascular dysfunction ( $LVEF \leq 30\%$ ). Sensitivity analysis as in the supplemental material of Lee and coworkers.<sup>48</sup>



### **BOX 1. Previous relevant studies on the use of levosimendan in cardiac surgery**

- Over 40 clinical trials were run on the use of levosimendan in cardiac surgery;
- Earlier studies suggested that levosimendan could prevent the development of LCOS and could be effective in treating postoperative LCOS;
- Meta-analyses suggested a reduction of mortality, significant when levosimendan is used in case of severe perioperative cardiovascular dysfunction ( $LVEF \leq 30\%$ );
- Indications of favorable renal effects in this setting have also been reported.

### **BOX 2. The three most recent studies on levosimendan in cardiac surgery**

- **LICORN** - Levosimendan on Low Cardiac Output Syndrome in Patients With Low Ejection Fraction Undergoing Coronary Artery Bypass Grafting With Cardiopulmonary Bypass trial (NCT02184819) assessing the efficacy of a pre-operative infusion of levosimendan in reducing postoperative LCOS in patients with poor LVEF undergoing CABG;
- **CHEETAH** - Levosimendan to Reduce Mortality in High Risk Cardiac Surgery Patients trial (NCT00994825) assessing the effect of levosimendan on cardiac surgery patients who developed postoperative LCOS;
- **LEVO-CTS** - Levosimendan in Patients with Left Ventricular Systolic Dysfunction Undergoing Cardiac Surgery Requiring Cardiopulmonary Bypass, Phase III clinical trial (NCT02025621) assessing the effect of levosimendan on patients with low pre-operative LVEF ( $EF < 35\%$ ) undergoing scheduled or urgent cardiac surgery.

**BOX 3. Consensus on efficacy and safety of levosimendan in operative settings**

Taking all the available data into consideration, including the experience of these three recent studies:

- levosimendan is a safe agent for the treatment of patients undergoing cardiac surgery and in need of inotropic support, despite the three large multi-center trials could not rule out their null hypothesis;
- levosimendan is an effective agent as it regards hemodynamic support;
- statistically significant mortality benefit seems to be limited to sub-groups, such as the isolated CABG procedures, and the low EF patients;
- further in-depth assessment of the utility of levosimendan will require additional trials in closely defined patient populations with study designs that mitigate to the fullest extent possible any influence of methodological variation.

**Table I: Studies on the use of levosimendan in peri-operative settings included in the meta-analysis by Harrison et al.<sup>17</sup>**

first author, year of publication (ref)	Settings	Trial design	LS dose	Start of treatment	Comparator	Patients in LS arm vs comparator	Primary end-points/evaluations	Secondary end-points/evaluations
Al-Shawaf 2006. <sup>20</sup>	Postoperative LCOS (within 12 h) in pts with type 2 DM and preoperative LVEF < 35%, to whom CABG was performed	Randomized open-label	12 mcg/kg bolus + 0.1-0.2 mcg/kg/min for 24 h	Within 12 h post surgery	Milrinone 50 mcg/kg bolus + 0.3-0.5 mcg/kg/min for 24 h	14 vs. 16	Superior hemodynamic effects (cardiac index, mixed venous saturation) with LS	No difference in insulin requirements, Time in ventilator & ICU shorter with LS
Alvarez 2005. <sup>21</sup>	Post-operative cardiac index < 2.5 l/min/m <sup>2</sup> in pts to whom CABG was performed	Randomized open-label	12 mcg/kg bolus + 0.2 mcg/kg/min for 24 h	Post surgery	Dobutamine 7.5 mcg/kg/min for 24 h	15 vs. 15	Cardiac index and heart rate increased, and mean arterial pressure, systemic and pulmonary vascular index decreased significantly in LS group	
Alvarez 2006. <sup>22</sup>	Postoperative LCOS (within 4 h) in pts to whom CABG or valvular surgery was performed	Randomized open-label	12 mcg/kg bolus + 0.2 mcg/kg/min for 24 h	Within 4 h post surgery	Dobutamine 7.5 mcg/kg/min for 24 h	25 vs. 25	More pronounced heart rate and cardiac index increase, and more pronounced decrease in systemic vascular resistance in LS group	The hemodynamic effects lasted beyond study drug infusion period only in LS group
DeHert 2007. <sup>23</sup>	Patients with preoperative LVEF ≤30% and to whom CABG and or valvular surgery was performed	Randomized open-label; assessment of outcomes by blinded observers	0.1 mcg/kg/min for 19 h	Immediately after release of aortic crossclamp	Milrinone 0.5 mcg/kg/min for 83 h	15 vs. 15	Stroke volume increased initially similarly, but the effect lasted longer in LS group	Secondary inotrope and vasoconstrictor need lower in LS group Tracheal intubation time shorter in LS group
Eriksson 2009. <sup>24</sup>	Patients with 3-vessel coronary disease and LVEF < 50% to whom CABG was performed	Randomized double-blind	12 mcg/kg bolus + 0.2 mcg/kg/min for 24 h	Immediately after induction of anesthesia	Placebo	30 vs. 30	Primary weaning from cardiopulmonary bypass successful in 73% vs. 33% in LS and placebo groups, respectively; p=0.002	Need for additional inotropic or mechanical therapy lower in LS group
Levin 2009. <sup>25</sup>	Postoperative LCOS (within 6 h) in pts with preoperative LVEF < 25%, to whom CABG was performed	Randomized open-label	10 mcg/kg bolus + 0.1 mcg/kg/min for 24 h	Within 6 h post surgery	Dobutamine 5-12.5 mcg/kg/min for 24 h	127 vs. 126	Post-operative mortality lower in LS group (7.1% vs. 15.9%; p<0.05)	Lower need for secondary inotropes (14.2 vs. 32.5%), vasopressors (17.3 vs. 43.6%) and intra-aortic balloon pump (3.1 vs. 14.2%); p<0.05 for all
Levin 2012. <sup>26</sup>	Patients with preoperative LVEF <25% undergoing CABG	Randomized open-label	10 mcg/kg bolus + 0.1 mcg/kg/min for 24 h	24 h before surgery	Placebo	127 vs. 125	Postoperative LCOS (7.1% vs. 20.8%; p<0.05) and mortality (3.9% vs. 12.8%; p<0.05) lower in LS group	Difficult weaning from cardiopulmonary bypass, need for secondary inotropes and vasopressors and need for intra-aortic

								ballon pump less frequent in LS group
Lomivorotov 2011. <sup>27</sup>	40 consecutive patients with LVEF<35%, who underwent coronary artery bypass grafting, were included	Randomized open-label	12 mcg/kg bolus + 0.1 mcg/kg/min for 24 h	Immediately after induction of anesthesia	Intra-aortic balloon pump 16-18 h before operation	20 vs. 20	Cardiac index significantly higher 5 min after CBP, at the end of the operation, 2 and 4 h after perfusion in LS group	Toponin I lower at 6 h after the operation and ICU stay in LS group
Järvelä 2008. <sup>28</sup>	Patients undergoing aortic valve surgery with or without CABG	Randomized double-blind	0.2 mcg/kg/min for 24 h	Immediately after induction of anesthesia	Placebo	12 vs. 12	LVEF maintained in LS group, but decreased in control group after operation	LS prevented postoperative decreases in mixed venous saturation and central venous oxygen saturation
Lahtinen 2011. <sup>29</sup>	Patients undergoing CABG or valve operation or both	Randomized double-blind	24 mcg/kg bolus + 0.2 mcg/kg/min for 24 h	Immediately after induction of anesthesia	Placebo	99 vs. 101	Heart failure defined as cardiac index <2.0 l/min/m <sup>2</sup> or failure to wean from CPB 15% vs. 58% in LS and placebo groups, respectively; p<0.001	Rescue inotrope and intra-aortic balloon pump needed less, but vasopressor more in LS group
Leppikangas 2011. <sup>30</sup>	Patients undergoing aortic valve surgery and CABG; additionally preoperative LVEF<50% or LV thickness > 12 mm	Randomized double-blind	12 mcg/kg bolus + 0.2 mcg/kg/min for 24 h	24 h before surgery	Placebo	12 vs. 12	Cardiac index and stroke volume index (SI) higher in LS group) for the 4 day postoperative period	
Momeni 2011. <sup>31</sup>	Pediatric study on patients between 0 and 5 y.o. requiring inotropic support for corrective congenital heart surgery under cardio-pulmonary bypass	Randomized double-blind	0.05 mcg/kg/min	onset of CPB	Milrinone	18 vs 18	No significant difference between serum lactate levels of groups.	Rate-pressure index significantly lower in LS group at 24 and 48 h, indicating lower myocardial oxygen demand
Tritapepe 2006. <sup>32</sup>	Patients undergoing elective CABG	Randomized double-blind	24 mcg/kg as a bolus in 10 min	Just before placing patient on CPB	Placebo	12 vs. 12	Lower troponin I release (p<0.05) and a higher cardiac index (p<0.05) postoperatively	
Tritapepe 2009. <sup>33</sup>	Patients undergoing elective CABG	Randomized double-blind	24 mcg/kg as a bolus in 10 min	Just before placing patient on CPB	Placebo	52 vs. 50	Length of ICU stay shorter in LS group: 25(7) vs. 32 (13) hours; p=0.002	Length of hospital stay and tracheal intubation time shorter in LS group. Troponin I release lower in LS group





